

Octopus consumption-based estimates

- •Octopus consumption by Bering Sea P. cod accepted as basis for recommendation in 2011; agreed to update every 5 years.
- Method performed for GOA octopus; GOA team preferred survey biomass (survey biomass currently used in GOA).







Why for Octopus, but not squid?

- Same species in diets and fishery.
- Same locations/depths.
- P. cod (well-monitored in biomass and diets) large proportion of octopus consumption. Squid consumed by many predators (including birds, mammals).
- Disadvantage for both: consumption is of small/juveniles, fishery is on adults.

Consumption Estimate: Method

M* B (cod eating octopus, t/year) =

Cod Ration_{length} (g/fish/year) *


Cod Nsurvey_{length,strata,year} (fish) *

Ration @ cod length: annual ration, calculated by von Bertalanffy growth fitting (Holsman and Aydin 2016). Non time-varying

Survey N (including uncertainty), more conservative than stock assessment, not dependent on stock assessment choices (CIE suggestion: try both).

Diet Comp survey length bin, strata, year (proportion)







Diet composition

- Calculated per-strata and in three broad length categories.
- Changes from last assessment
 - Adjusted length bins to better extrapolate missing data
 - Moved from direct empirical method to fitting Dirichlet distribution
 - Several more years of data



Estimated Octopus Consumption (old)








OFL: Harmonic mean of harmonic means: 3,452 t (old) 4,770 t (new)

Bering Catch 500t (TAC-limited); GOA going up (900t)

Estimated Octopus Consumption (new)







Questions

- Use all years, accept jump?
- Use new years in future?
- Add 2012/2013 (and 2016) next year or in 5 years?
- Other methodological concerns?
- Other requested explorations or analyses?

