

Operating Model for Halibut ABM

SSC Recommendations - June 2018


Based on these criteria, the SSC recommends that the working group move forward with their proposed plan for the October 2018 meeting using Approach 2 (i.e., the “two-area” halibut simulation model) to conduct a preliminary analysis of a subset of ABM alternatives. Approach 2 is the simplest and the closest to being developed of the three suggested approaches and will be capable of providing a preliminary analysis of ABM alternatives for the October 2018 meeting. Further, this approach is the most transparent and can explore some of the key assumptions that underlie halibut population dynamics. However, in its current form, Approach 2 still requires additional work before it can be used to evaluate ABM alternatives.

SSC Recommendations - June 2018 cont.

- **A management model for setting the Total Constant Exploitation Yield (TCEY) must be developed.** The SSC agrees with the working group's suggested approach of using historical observations to relate available halibut biomass and TCEY.
- **Halibut PSC limits must be tied to impacts in the groundfish and directed halibut fisheries.** The SSC was unsure of whether the suggested empirical approach of relating historical PSC limits to PSC use and groundfish harvest is appropriate since there is very little historical variation in PSC limits. A different approach for measuring impacts to the groundfish fisheries may be needed.
- **Additional thought must be given to how uncertainty will be incorporated into the model.** The working paper suggests that a halibut simulation framework with "perfect information" could be ready for the October 2018 meeting, but the SSC notes the importance of evaluating the performance of ABM alternatives under various sources of uncertainty (e.g., estimation, process, or implementation error).


Operating Model (OM) Overview

- Sex and age-structured
- 2 Areas
 - BSAI region (4ABCDE)
 - Remaining regions in aggregate
- Common recruitment
 - Apportioned among areas
- 3 Gear types (Selectivity and F_t)
 - Directed Fishery, PSC Trawl, PSC Longline
- Gear-specific Selectivity
- Age-specific movement between areas




Operating Model (OM) Overview

- Sex and age-structured
- 2 Areas
 - BSAI region (4ABCDE)
 - Remaining regions in aggregate
- Common recruitment
 - Apportioned among areas
- 3 Gear types (Selectivity and F_t)
 - Directed Fishery, PSC Trawl, PSC Longline
- Gear-specific Selectivity
- Age-specific movement between areas




Operating Model (OM) Overview

- Sex and age-structured
- 2 Areas
 - BSAI region (4ABCDE)
 - Remaining regions in aggregate
- Common recruitment
 - Apportioned among areas
- 3 Gear types (Selectivity and F_t)
 - Directed Fishery, PSC Trawl, PSC Longline
- Gear-specific Selectivity
- Age-specific movement between areas




Operating Model (OM) Overview

- Sex and age-structured
- 2 Areas
 - BSAI region (4ABCDE)
 - Remaining regions in aggregate
- Common recruitment
 - Apportioned among areas
- 3 Gear types (Selectivity and F_t)
 - Directed Fishery, PSC Trawl, PSC Longline
- Gear-specific Selectivity
- Age-specific movement between areas



Movement Among Areas



Movement Among Areas



Survival

Area: BSAI

Area: Other


Years →
← Earlier

Later




Movement Among Areas

Survival
Movement **Out** of BSAI



Movement Among Areas

Survival
Movement **Out** of BSAI
Movement **Into** BSAI



Sources of Operating Model ***Variation***

- Current
 - Log-normal process variation in recruitment
- Potential
 - Temporal variation in weight at age
 - Temporal variation in mean recruitment (periodic, PDO-like)

Operating model major uncertainties

- Recruitment distribution between GOA and BS
- Migration rates
- Weight-at-age regime
- Recruitment regime (e.g. high or low period for R_0)

Operating model major uncertainties

- Recruitment distribution between other regions and BSAI
- Migration rates
- Weight-at-age regime
- Recruitment regime (e.g. high or low period for R_0)

Recruitment allocation

IPHC research on larval distribution (past and current)


<https://iphc.int/management/science-and-research/biological-and-ecosystem-science-research-program-bandesrp/bandesrp-migration/larval-distribution>

[IPHC-2018-SRB012-09](#), page14

- Prevailing currents transport larvae from east to west
- NOAA ichthyoplankton surveys in GOA and BS
 - 1972-2015
 - Larval flow through Unimak Pass
- Space-time modelling of Pacific halibut larval observations
 - Comparison of distribution in warm years (2001-2005) and cold years (2007-2013)
- Collaboration with NOAA/EcoFOCI
 - Larval transport model to examine currents and larval advection


Movement data and rates

- Current knowledge of migration ([IPHC-2018-AM094-09](#), pages 46-47)
- Adult movement rates reasonably well understood
 - Historical and recent PIT tagging studies
- Less than age 5, trawl gear tagging studies
- Less data from IPHC Reg Area 4
 - Appreciable movement out of Area 4 between ages 5-10
 - Paucity of data for ages 2-4
- Assumptions
 - Ages 0-1 do not move
 - Age 2 cannot move from Area 4 to Area 2
 - Ages 2-4 in/out of Area 4 based relative estimates of migration from Area 3 to Area 2



Movement from IPHC data

- Probability of movement determined from IPHC estimates of Area 4 to all other areas
 - All Areas to Area 4 is zero across all ages
- Extremely high recruitment necessary in BSAI to maintain halibut in the BSAI
- Research is being conducted on migration
 - [IPHC-2018-SRB012-09](#), page15



Key Operating Model *Uncertainties*

The Outflow Problem

- Recruitment allocation among areas – this example shows all recruitment to BSAI
- Model has age-specific movement rates among areas, but these examples show movement constant across ages and sexes





Figure 3. Simulated female spawning stock biomass at age over time under two movement scenarios, with no fishing mortality and all recruitment allocated to the BSAI area. Annual movement rates are constant across ages and sexes.

Recruitment Allocation vs. Movement Rate

Equilibrium relative biomass at age in each area


- The combination of recruitment allocation and annual movement rate dictates both the equilibrium biomass in each region and the age-specific contribution to total biomass



Recruitment Allocation vs. Movement Rate

Equilibrium relative biomass at age in each area

- The combination of recruitment allocation and annual movement rate dictates both the equilibrium biomass in each region and the age-specific contribution to total biomass



Future Work Plan

- Identify (realistic and stable) trial values for parameters describing
 - Recruitment apportionment
 - Movement among areas
- Quantify TCEY -> FCEY pathway and harvest allocation procedure
- Add estimation (model) uncertainty to the management process
- Define selectivity for all gears (within areas)
- Add weight-at-age scenarios
 - Through variation in weight-at-age vector
- Add recruitment regime scenarios or periodic regime shifts
- Use sensitivity scenarios (high, low) for growth and recruitment or time-varying simulation?

Supplementary Slides


Recruitment: Beverton-Holt with Steepness $h=0.75$

Recruitment by Area and Year

$$R_{l,y} = \delta_l \frac{SSB_y 4hR_0}{SSB_0(1-h) + SSB_y(5h-1)} e^{\varepsilon_y - \frac{\sigma_r^2}{2}}$$

Recruitment Process Error

$$\varepsilon_y \sim \text{Normal}(0, \sigma_r)$$



Recruitment: Beverton-Holt with Steepness $h=0.75$

Recruitment by Area and Year

$$R_{l,y} = \delta_l \frac{SSB_y 4hR_0}{SSB_0(1-h) + SSB_y(5h-1)} e^{\varepsilon_y - \frac{\sigma_r^2}{2}}$$

Annotations for the equation:


- δ_l : Proportion of Recruitment by area
- SSB_0 : Equilibrium Spawning Stock Biomass
- R_0 : Equilibrium Recruitment

Recruitment Process Error

$$\varepsilon_y \sim \text{Normal}(0, \sigma_r)$$

Annotations for the equation:

- σ_r : Process Error Standard Deviation: 0.6



Recruitment: Spawning Stock Biomass Calculation

- Product of female biomass at age and maturity at age, summed across areas and ages

$$SSB_y = \sum_l \sum_a N_{l,s,y,a} w_{s,a} m_{s,a} = \sum_l \sum_a B_{l,s,y,a} m_{s,a}$$

Abundance by area, sex, year, and age

Weight by sex and age

Maturity by sex and age
Note: $m=0$ for males at all ages

Biomass by area, sex, year, and age

- Recruitment at to age-1 assumes a 50/50 sex ratio

$$N_{l,s,y,a=1} = 0.5 R_{l,y-1}$$

Recruitment from previous year

Updating Numbers (Abundance) at Age

Total Instantaneous Mortality

$$Z_{l,s,y,a} = M_s + \sum_g v_{g,s,a} F_{l,g,y}$$

Natural Mortality by sex

Fishing Mortality by area, gear, and year

Selectivity by gear, sex, and age

Numbers at age for all ages $1 < a < A$ (plus age group A: 30)

$$N_{l,s,y,a} = N_{l,s,y-1,a-1} e^{-Z_{l,s,y-1,a-1}}$$

Updating Numbers at Age: Plus Age Group

Plus age group $A=30$

$$N_{l,s,y,a=A} = N_{l,s,y-1,a=A} e^{-Z_{l,s,y-1,a=A}} + N_{l,s,y-1,a-1} e^{-Z_{l,s,y-1,a-1}}$$

Surviving Individuals in
Plus Age Group

New Entrants to Plus
Age Group

Harvest Calculations

Age-specific total catch in numbers by year

$$C_{l,s,y,a} = \left(\frac{f_{l,s,y,a}}{Z_{l,s,y,a}} \right) N_{l,s,y,a} \left(1 - e^{-Z_{l,s,y,a}} \right)$$

Total Instantaneous Mortality

Gear-specific annual catch

$$c_{l,s,y,a,g} = \left(\frac{v_{g,s,a} F_{l,g,y}}{Z_{l,s,y,a}} \right) N_{l,s,y,a} \left(1 - e^{-f_{l,s,y,a}} \right)$$

Sum of Fishing Mortality
Across Gear Types

$$f_{l,s,y,a} = \sum_g v_{g,s,a} F_{l,g,y}$$

Gear-specific annual harvest in units of biomass

$$H_{l,y,g} = \sum_s \sum_a c_{l,s,y,a,g} w_{s,a}$$

Movement Among Areas

- Movement *may* be age-specific
- Occurs after natural and fishing mortality, but prior to SSB calculations

Number of migrants from area i to area j

$$\tau_{i,j,s,y,a} = N_{l=i,s,y,a} \pi_{i,j,a}$$

Transition Probability at age

Updating abundance by area, sex, year, and age

$$N_{l,s,y,a} = N_{l,s,y,a} + \sum_{k \in \text{areas}} \tau_{i=k,j=l,s,y,a} - \sum_{k \in \text{areas}} \tau_{i=l,j=k,s,y,a}$$

Immigrants **into** area a Emigrants **out** of area a

Notation: Symbols

Symbol	Description
<i>l</i>	Area or location (Bering Sea and Aleutian Islands and remaining West Coast halibut range)
<i>y</i>	Year
<i>s</i>	Sex
<i>a</i>	Age
<i>g</i>	Gear type or fishing sector
<i>i</i>	Area migrating from
<i>j</i>	Area migrating to

Notation: Derived Parameters

Parameter	Description
$R_{l,y}$	Recruitment
SSB_y	Spawning stock biomass
$N_{l,s,y,a}$	Numbers at age
$B_{l,s,y,a}$	Biomass at age
$Z_{l,s,y,a}$	Total mortality
$F_{l,g,y}$	Fishing mortality rate
$f_{l,s,y,a}$	Age and sex-specific fishing mortality rate
$C_{l,s,y,a}$	Total catch in numbers
$c_{l,s,y,a,g}$	Catch in numbers by gear type
$H_{l,y,g}$	Harvest in biomass by gear type

Notation: Input Parameters

Parameter	Description
M_s	Natural mortality by sex
$w_{s,a}$	Weight at age by sex
$m_{s,a}$	Maturity at age (note this is equal to zero for males)
$v_{g,s,a}$	Selectivity
B_{start}	Initial biomass
$p_{l,s,a}$	Initial biomass proportions at age by area

Operating Model (OM) Overview

- Sex and age-structured
- 2 Areas
 - BSAI region (4ABCDE)
 - Remaining West Coast distribution
- Common recruitment
 - Apportioned among areas
- 3 Gear types (Selectivity and F)
 - Directed Fishery, PSC Trawl, PSC Longline
- Gear-specific Selectivity
- Age-specific movement between areas

